



Effects of *Rosmarinus Officinalis* Plant Extract on *Trichomonas Vaginalis* Parasites and *Candida albicans* under Laboratory Conditions: An Experimental Study

Saeide Saeidi ^{1,*}, Forogh Forgani ², Fereshteh Javadian ² and Elham Javadian ^{3,**}

¹Zabol University, Zabol, Iran

²Zabol University of Medical Sciences, Zabol, Iran

³Zahedan University of Medical Science, Zahedan, Iran

*Corresponding author: Zabol University, Zabol, Iran. Email: s.saeedi2@yahoo.com

**Corresponding author: Zahedan University of Medical Science, Zahedan, Iran. Email: elham.javadian1366@gmail.com

Received 2019 May 04; Revised 2019 July 08; Accepted 2019 July 17.

Abstract

Background: As we know metronidazole is a selective treatment for Trichomoniasis, however, it has many side effects, and, on the other hand, it is important to try finding natural alternative treatments.

Objectives: The aim of this study was to investigate the effects of *Rosmarinus officinalis* plant extract on *Trichomonas vaginalis* parasites and *Candida albicans* under laboratory conditions.

Methods: In this experimental study, *Rosmarinus officinalis* methanolic extracts were prepared. The anti-trichomoniasis effect was evaluated at concentrations of 0.1, 0.01, 0.001, 0.0004, 0.0002, and 0.0001 $\mu\text{g}/\text{mL}$ in TYIS33 culture media and resulted data were analyzed by ANOVA.

Results: The results of this study showed that in the concentration of 0.001 at 4 hours, the parasite was spherical and inactivated, while at concentrations of 0.0004, 0.001, 0.01, and 0.1 in 1 hour. The results of this study showed that *Rosmarinus officinalis* extract in concentration of 100 $\mu\text{g}/\text{mL}$ has an inhibitory effect on *Candida albicans*, so that the highest inhibitory diameter was 18.3 ± 0.9 mm while two other strains exhibit resistance and no inhibitory zones.

Conclusions: The results of this study showed that rosemary extract inhibits the growth of *Trichomonas vaginalis* and *Candida albicans* that can be used to treat infections.

Keywords: *Trichomonas Vaginalis*, *Candida albicans*, Rosemary, Laboratory Conditions

1. Background

Vaginitis is one of the most common diseases of women, which is important in health centers around the world and requires health care (1).

About 10 million health centers annually address vaginal discharge complaints. Vaginal symptoms are usually related to bacterial vaginitis, vulvovaginal candidiasis, and trichomoniasis, in addition, Chlamydia trachomatis and Nisria gonorrhoea are other causes of vaginitis (2).

Trichomonas vaginalis is a flagellate protozoan transmitted through sexual contact (3). It is also the only pathogenic form of the parasite in humans, this parasite grows in a humid environment, pH ranges from 4.9 to 7.5 and temperatures ranging from 35°C to 37°C. If these conditions turn less or more than optimal, the organism disappears. Women usually have vaginal and urethral infection, however, the parasite may also infect the cervix, the

Bartholin glands, or the bladder. In men, the organism is often found in the extremities of the urethra, however, it involves the prostate gland, seminal sac, and epididymis rarely.

The parasite causes various diseases like vaginitis, urethritis, and prostatitis (4).

This parasite has a 15% - 10% prevalence and is one of the most common sexually transmitted diseases after bacterial infection of Chlamydia. With the report of the World Health Organization, 170 - 190 million people worldwide are now infected with this parasite, with 120 million cases of trichomoniasis annually, by checking the results; we find that nearly 20% of cases show vaginitis symptoms.

According to a recent systematic study, the prevalence of trichomoniasis in Iran is estimated at 8% (5).

Prevalence increases by up to 30% in high risk populations (6). It should be noted that pregnant women with

trichomoniasis may show multiple side effects, including preterm labor, low birth weight, premature rupture of the membrane, tubal infertility, and atypical pelvic inflammatory disease. On the other hand, increasing the transmission risk of HIV/AIDS, human papillomavirus, and cervical cancer are another serious complication of this disease (7). Men with *Trichomonas vaginalis* are usually unmarked and show symptomatic infection that is associated with urethritis (urticaria and secretion) (4). Trichomoniasis in men is usually treated without medication (8). This wide spread use of herbal medicine can be due to various reasons, such as fewer side effects, better patient acceptance due to traditional medicine advice, lower prices for medicinal plants, and compatibility with the physiological function of the human body. Herbal medicine is a branch of traditional medicine in ancient civilizations like Iran, which played a major role in the treatment of diseases one century ago.

Candidiasis is one of the most common opportunistic fungal diseases in humans. This disease caused by the yeast fungus is called candida (9). *Candida albicans* yeasts are commonly found in humans, and their growth is typically limited by the human immune system and other microorganisms, such as bacteria that occupy identical places in the human body. Candida infection shows a range of surface complications such as oral thrush and vaginal inflammation that potentially treats by itself in humans (10).

Candida's genus contains a heterogeneous group of organisms that grows in yeast, and most members of this genus produce false filaments during their growth; however, *Candida albicans* and *Candida Doblini Ninci* have the true form of Heif and cells with thick walls; both of them produce Chlamydosporum, which are detectable in a laboratory.

Candida species are the fourth cause of blood infections in hospitalized patients and they can potentially cause the mortality of patients that have been admitted to hospitals in the United States (11).

However, there are also limitations in the treatment of fungal diseases, such as the prevalence of resistant species to the antifungal agents and their side effects. Therefore, studying on new drugs that have fewer side effects for patients at the same time with eliminating or inhibiting the pathogen is necessary. For this suggestion, studying on medicinal plants is important.

Rosemary (*Rosmarinus officinalis* L.) is a plant from the family of mint, which is native to the Mediterranean area, and also in Iran.

2. Objectives

We are familiar with anticancer, anti-inflammatory, antimicrobial, and antioxidant effects of this herb (12). The main purpose of this study is to investigate the effects of methanolic extract of *Rosmarinus officinalis* on *Trichomonas vaginalis* and *Candida albicans* under laboratory conditions.

3. Methods

3.1. Extract Preparation

The massage method was used to prepare extracts. In this way, after crushing the leaves of the plant, 50 g of each sample was embedded in methanol for 48 hours, and after this time a filter is used for material separation then, the extracts were concentrated by using a rotary machine (vacuum distillation) at 40°C to 50°C, and finally, dried at 40°C for two days.

The *Rosmarinus officinalis* plant used in this research was collected from the city of Zabol and was diagnosed by Rosemary officinalis by the expert of Zabol National University, according to the visual characteristics and herbalists' descriptions. After collecting the leaves, the leaves were dried in the shade in order to prepare the extract with crushed mill. Then, vaginal discharge of women with vaginitis symptoms was observed directly in the treatment centers. After confirmation, *Trichomonas* parasite was cultured in a media and prepared eggs. Five tubes containing culture media and *T. vaginalis* were considered as controls. Five tubes containing metronidazole and *T. vaginalis*, DMSO and *T. vaginalis*, and dry extract at concentrations of 0.1, 0.01, 0.001, 0.0004, 0.0002, and 0.0001 mg/mL in DMSO solvent and *T. vaginalis* were prepared in a 37-degree incubator.

3.2. Isolation of *Candida albicans*

Samples were taken by two sterile swabs of vaginal discharge. Samples were taken in pipettes containing 2 mL of sterilized distilled water while transferring to a laboratory. The swabs were used for direct testing with 15% potassium, cultured on Sabouraud dextrose agar (Merck, Germany) and chromogen agar (France, France). The culture media was incubated for 48 to 72 hours at 30 degrees Celsius to confirm the diagnosis of the albicans species from other species.

Candida from the colonies on the corn-mili agar + tween media linear cultures was given. In addition, tuberculosis and beta-glucosidase tests were used to differentiate *Candida albicans* from other species. The plates were placed at 25°C for 72 hours. From a fresh culture, single

colonies of each yeast were transferred to a solution of 20% water-glycerol and the sample was transferred. They were kept at a minimum temperature of 20°C, the samples were stored in the solution of 20 g/g glycerol in case it was needed to repeat each step of the experiment.

3.3. Disc Diffusion Method Was Used for Determining the Diameter of the Extract Halo on *Candida albicans*

Agar diffusion method and disc diffusion method were also used to determine the antifungal activity of medicinal plants extract against opportunistic fungi (13). It should be noted that the agar diffusion method was subjected to minor changes (14).

In this method, according to the relevant standard, after accurate counting of fungi and preparation of standard suspensions with a certain amount of fungi, it was transferred into a media and cultured using sterilized swabs. Then, the wells with a diameter of 5 mm were placed on a subrock dextrose agar media, and the ends of each well were closed to a substrate in sterile conditions, which resulted in preventing the extract from being extracted below the culture media. Next, extract in these wells that have different concentrations of 2.5 - 40 µg/mL were sold and incubated at 37°C, after 24°C - 48°C. The non-growth halo diameter was investigated.

3.4. Determination of the Minimum Inhibitory Concentration of MIC Extract on *Candida albicans*

To determine the MIC for *Rosmarinus officinalis* extract separately, a series of 9 tubes are used to test the dilutions of each extract and a tube as a positive control containing diluted extract and culture media and negative control including microbial suspension and culture media. We considered that the initial concentration of the extract is 50 mg/mL, which is obtained by inserting 1 mL of the extract into the first tube containing one ml of culture media at a concentration of 25 µg/mL. In this way, for the first tube, one milliliter of the extract is diluted at a concentration of 100 µg/mL with one mL of the dextrose broth agar media, and, similarly, taking 1 mL of the first tube, and pouring it into the second tube containing one milliliter The litter is Dextrose Broth Agar, which is transmitted to the last tube. The tube was lifted out of one milliliter and poured out. By doing this process, the dilution of each tube was half the dilution of the tube before. A total of 50 µL of microbial suspension was transferred into all the tubes except the positive control tube. The dilutions of the extracts for our studied fungi were performed in a completely separate manner. The series of test tubes were placed in an incubator for 24 hours at 37°C, then, the tubes were examined for turbidity due to growth of the fungus.

The results were expressed as mean or ranked in order of importance as percent. The data were subjected to one-way analysis of variance (ANOVA), using the SPSS V. 17 software. The P value of less than 0.05 were regarded as significant.

4. Results

The results of this study showed that in the concentration of 0.0001 at 4 hours, the parasite Spherical and inactive, dead. At a concentration of 0.0002 in 2 hours, the parasite is spherical and inactivated (Table 1).

The results of this study showed that *Rosmarinus officinalis* extract in concentration of 100 µg/mL has an inhibitory effect on *Candida albicans*, therefore, the highest inhibitory diameter was 18.3 ± 0.9 mm while two other strains exhibit resistance and no inhibitory zones (Table 2).

5. Discussion

Trichomonas vaginalis acts as a predisposing factor in the transmission of human immunodeficiency virus and other sexually transmitted infections (15). According to numerous reports about drug resistance of *Trichomonas vaginalis* to metronidazole, which is the main treatment of this infection, it is needed to find an effective drug with minimum complications (16).

Candida albicans is an eukaryotic fungal pathogen that causes diseases like oral thrush and vulvovaginitis. Many of its biological pathways are common with humans, and most antifungal drugs have different side effects at different doses. Despite the effect of these drugs on the patient's recovery, excessive use of drugs caused drug resistance, which has led researchers to use plant compounds as an alternative for antibiotics to inhibit the growth of these pathogens in research's.

Rosmarinus officinalis is used as a styptic agent, antipyretic, antispasmodic, and sudorific in traditional medicine. Extract and volatile oil are used for abortion and to increase menstrual bleeding. The use of rosemary extract in cosmetics is very common, and evidence suggests the effect of its lotion on stimulating hair growth and prevention of scabies. Historical reports of rosemary treatment are available as herbal medicine. Rosemary is one of the oldest medicinal plants known to have been used centuries ago to strengthen memory and brain activity.

Major components of the volatile oil include: Camphor, Borneol, α-Pinen, the amount and percentage of each of these materials depends on the environmental conditions of plant's site.

Table 1. Effect of Different Concentrations of *Rosmarinus officinalis* Extract on *Trichomonas Vaginalis* Parasite

Extract Concentration	At the Time of Cultivation	Hour 1	Hour 2	Hour 4
0.1	Dead and without walls	-	-	-
0.01	Dead and without walls	-	-	-
0.001	Spherical and inactive	-	-	-
0.0004	Live with low mobility and dead	-	-	-
0.0002	+	Live with low mobility	Spherical and inactive	-
0.0001	+	+	Live with low mobility	Spherical and inactive, dead

Table 2. Inhibitory Zones Extract Concentration 100 µg/mL Against *C. albicans* (mm)

Strain Fungal <i>C. albicans</i>	Inhibitory Zones	Strain Fungal	Inhibitory Zones
1	9.2 ± 0	7	17.4 ± 0.5
2	18.3 ± 0.9	8	-
3	9.5 ± 0.3	9	9.6 ± 0.5
4	10.2 ± 0.5	10	-
5	14.6 ± 0.8	11	15.3 ± 0.5
6	12.5 ± 0.6	12	12 ± 0.8

Other ingredients in Rosemary leaf include: luteonolin, jenkannin, tannin, resin, pasonin, fat, carbohydrates, and vitamins.

The results of this study showed that *Rosmarinus officinalis* has good anti-fungal and anti-parasitic activities.

In another study, the results showed that *Rosmarinus officinalis* essential oil inhibits bacteria including *E. coli*, *Bacillus cereus*, *Staphylococcus aureus* (17), *Clostridium perfringens*, *Aeromonas hydrophila*, *Bacillus cereus*, and *Salmonella choleraesuis*.

Essential oils of *Zataria multiflora* and *Myrtus communis* showed inhibitory effects at concentrations of 0.1%, 0.01%, 0.001%, and 0.004% on *Trichomonas vaginalis* (18). Meanwhile the methanolic extract of *Myrtus communis* showed an inhibitory effect at concentration of 0.1 mg/mL (19).

In another study, aqueous extracts was also tested, *Verbena* sp. (*Guachu ka'a* in Mbya-Guarani language) and *Campomanesia xanthocarpa* (*Guavira* in Mbya-Guarani language) showed that the highest activity against *T. vaginalis* with MIC value of 4.0 mg/mL reached 100% of efficacy against the parasite. The kinetic growth assays showed that the extracts promoted complete growth inhibition after 4 hours of incubation (20).

Plant of *Verbena* sp, *C. xanthocarpa*, *Myrtus communis* were tested against *T. vaginalis*, however, without success (21, 22).

In the study of Hassani et al., the effect of hydroalcoholic extract of eucalyptus in comparison with metronida-

zole on *Trichomonas vaginalis* has been evaluated under laboratory conditions. Extract of *E. camaldulensis* showed 80% growth inhibition (GI) in a concentration of 12.5 mg/mL during 24 hours. Diethyl ether extract in a concentration of 25 mg/mL showed 100% GI during 24 hours. With ethyl acetate extract, 100% GI was detected with the minimum concentration of 12.5 mg/mL in the first 24 hours. Finally, water extract in a concentration of 50 mg/mL showed 80% and 100% GI after 48 and 72 hours, respectively (23).

Amaryllidaceae species showed a promising activity against *Trichomonas vaginalis* (24, 25).

Also, different parts of the plants (leaves, flowers, etc.) also have antimicrobial properties (26).

In a study by Nejati et al., on 75 Wistar male rats (weighing 210 ± 10 g), after general anesthesia, a 1.5 to 1.5 cm squared sore wound made and immediately implanted and infected by *Candida albicans*. Experimental rats were randomly assigned into three groups of 25 (control and groups treated with 1.5 and 3% oat) and each group was divided into 5 subgroups of 5 (sampling groups in different days). During the course of the project, at the end of days 4, 8, 12, 16, and 20, wounds were collected from different groups for pathological examination by a special punch for biopsy. The results showed that application of 1.5% and 3% ointment from rosemary essential oil significantly reduced the infection rate and increased collagen content and production of coated tissue compared with the control group. Based on the results, the wound healing process in the ointment was 3% lower than the lower dose and the control showed a better result (27).

In the Natanzian Ghahfarkhi study, the antifungal effects of essential oils and alcoholic beverages on isolates that are clinically resistant and susceptible to fluconazole *Candida albicans* were examined *in vitro*, the inhibitory effect of alcoholic and oleaginous extracts of herbs on the growth of susceptible and resistant strains to fluconazole *Candida albicans* was confirmed (28).

In the study of Jafari et al., the concentration of 8.75 mg/mL from the extract of *Angus* spp. completely prevented the growth of *Candida albicans* and destroyed all the

live *Candida* cells at this concentration (MFC). In addition, concentrations of 4.23 mg/mL and 4.4 mg/mL aqueous extract were obtained as minimum inhibitory values of 50% and 90% candidiasis, respectively. In terms of fluconazole, 128 μ g/mL concentration was obtained as MFC and a concentration of 0.5 μ g/mL as MIC₅₀ (29).

In the study of Doddanna et al., which investigated the inhibition effect of the extract of the plant against *Candida albicans*, the results showed that the other plant extracts like alcoholic onion leaves, alcoholic tea leaves, alcoholic onion bulb, alcoholic Aloevera, and alcoholic mint leaves also inhibit the growth of *Candida albicans* but lesser than the extent (30).

In Prabhakar et al., which isolated 46 species of *Candida albicans*, antifungal activity of ethanol extracts of five plant species that included *Syzygium jambolanum*, *Cassia siamea*, *Odina wodier*, *Momordica charantia*, and *Melia azedarach* and two algal species, *Sargassum wightii* and *Caulerpa scalpelliformis* were tested against 25 isolated strains by disc diffusion method. Antifungal activity was observed at 100 mg/mL for *Syzygium jambolanum*, *Cassia siamea*, and *Caulerpa scalpelliformis*, and at 10 mg/mL for *Sargassum wightii* (31).

In another study, antifungal activity of *Mentha piperita*|*Mentha Arvensis*, *Aloe vera*, *Murraya koenigii*, *Allium cepa*, *Camellia sinensis* were investigated. The results showed that the raw extract of tea leaf had the highest inhibition zone against *Candida albicans* (30). In the study of Masom et al., which included antifungal activity of 5 plant species *Trachyspermum ammi* (seed), *Teucrium polium* (leaf), *Piper nigrum* (seed), *Pistachia vera* (skin), and *Camellia sinensis* (leaf), the results showed that all plant extracts inhibited *Candida albicans*, while the largest inhibition zone was for *Pistachio vera* (40 mm).

The lowest antifungal activity was produced by *Piper nigrum* (13 mm) and the minimum inhibitory concentration and the minimum inhibitory concentration of *Pistachio vera* were 6.25 and 12.5 mg/mL (32), respectively.

5.1. Conclusions

In general, since various studies have confirmed the antimicrobial, anti-cancer, and antioxidant properties of rosemary essential oil (33).

The results of this study showed that methanolic extract of rosemary at different concentrations and times has antifungal and antiparasitic effects that can be used to treat infections caused by them.

Footnotes

Authors' Contribution: All authors had an equal role in study design, work, statistical analysis, and manuscript

writing.

Conflict of Interests: The authors declare no conflict of interest.

Ethical Considerations: The code of ethics was zbmu.1.rec.1396.24.

Funding/Support: This study was supported by the Institute of Plant Biotechnology, University of Zabol, Zabol, Iran.

Patient Consent: The written informed consent was obtained.

References

- Johnson LF, Coetzee DJ, Dorrington RE. Sentinel surveillance of sexually transmitted infections in South Africa: A review. *Sex Transm Infect.* 2005;**81**(4):287-93. doi: [10.1136/sti.2004.013904](https://doi.org/10.1136/sti.2004.013904). [PubMed: [16061532](https://pubmed.ncbi.nlm.nih.gov/16061532/)]. [PubMed Central: [PMC1745020](https://pubmed.ncbi.nlm.nih.gov/PMC1745020/)].
- Aboud S, Msamanga G, Read JS, Mwatha A, Chen YQ, Potter D, et al. Genital tract infections among HIV-infected pregnant women in Malawi, Tanzania and Zambia. *Int J STD AIDS.* 2008;**19**(12):824-32. doi: [10.1258/ijsa.2008.008067](https://doi.org/10.1258/ijsa.2008.008067). [PubMed: [19050213](https://pubmed.ncbi.nlm.nih.gov/19050213/)]. [PubMed Central: [PMC2698963](https://pubmed.ncbi.nlm.nih.gov/PMC2698963/)].
- Kissing PJ, Dumestre J, Clark RA, Wenthold L, Mohammed H, Hagensee ME, et al. Vaginal swabs versus lavage for detection of *Trichomonas vaginalis* and bacterial vaginosis among HIV-positive women. *Sex Transm Dis.* 2005;**32**(4):227-30. [PubMed: [15788920](https://pubmed.ncbi.nlm.nih.gov/15788920/)].
- Petrin D, Delgaty K, Bhatt R, Garber G. Clinical and microbiological aspects of *Trichomonas vaginalis*. *Clin Microbiol Rev.* 1998;**11**(2):300-17. [PubMed: [9564565](https://pubmed.ncbi.nlm.nih.gov/9564565/)]. [PubMed Central: [PMC106834](https://pubmed.ncbi.nlm.nih.gov/PMC106834/)].
- Hezarjaribi HZ, Fakhari M, Shokri A, Teshnizi SH, Sadough A, Taghavi M. *Trichomonas vaginalis* infection among Iranian general population of women: A systematic review and meta-analysis. *Parasitol Res.* 2015;**114**(4):1291-300. doi: [10.1007/s00436-015-4393-3](https://doi.org/10.1007/s00436-015-4393-3). [PubMed: [25732256](https://pubmed.ncbi.nlm.nih.gov/25732256/)].
- Ziaei Hezarjaribi H, Dalimi A, Ghasemi M, Ghafari R, Esmaeili S, Armat S, et al. [Prevalence of common sexually transmitted diseases among women referring for pap smear in Sari, Iran]. *J Mazandaran Univ Med Sci.* 2013;**22**(1):19-24. Persian.
- Sood S, Kapil A. An update on *Trichomonas vaginalis*. *Indian J Sex Transm Dis.* 2008;**29**(1):7. doi: [10.4103/0253-7184.42707](https://doi.org/10.4103/0253-7184.42707).
- Weston TE, Nicol CS. Natural history of Trichomonal infection in males. *Br J Vener Dis.* 1963;**39**:251-7. doi: [10.1136/sti.39.4.251](https://doi.org/10.1136/sti.39.4.251). [PubMed: [14086515](https://pubmed.ncbi.nlm.nih.gov/14086515/)]. [PubMed Central: [PMC1047587](https://pubmed.ncbi.nlm.nih.gov/PMC1047587/)].
- Parks ET, Lancaster H. Oral manifestations of systemic disease. *Dermatol Clin.* 2003;**21**(1):171-82. viii. [PubMed: [12622279](https://pubmed.ncbi.nlm.nih.gov/12622279/)].
- Cortes JA, Concha MA, Cediel T, Castillo JS. [Diagnostic methods in candidemia: A systematic review of literature with meta-analysis]. *Rev Chilena Infectol.* 2011;**28**(5):423-8. Spanish. [PubMed: [22051618](https://pubmed.ncbi.nlm.nih.gov/22051618/)].
- Panacek A, Kolar M, Vecerova R, Pucek R, Soukupova J, Krystof V, et al. Antifungal activity of silver nanoparticles against *Candida* spp. *Biomaterials.* 2009;**30**(31):6333-40. doi: [10.1016/j.biomaterials.2009.07.065](https://doi.org/10.1016/j.biomaterials.2009.07.065). [PubMed: [19698988](https://pubmed.ncbi.nlm.nih.gov/19698988/)].
- Nieto G, Huvaere K, Skibsted LH. Antioxidant activity of rosemary and thyme by-products and synergism with added antioxidant in a liposome system. *Eur Food Res Tech.* 2011;**233**(1):11-8. doi: [10.1007/s00217-011-1486-9](https://doi.org/10.1007/s00217-011-1486-9).
- Duarte MC, Figueira GM, Sartoratto A, Rehder VL, Delarmelina C. Anti-*Candida* activity of Brazilian medicinal plants. *J Ethnopharmacol.* 2005;**97**(2):305-11. doi: [10.1016/j.jep.2004.11.016](https://doi.org/10.1016/j.jep.2004.11.016). [PubMed: [15707770](https://pubmed.ncbi.nlm.nih.gov/15707770/)].
- Jainkittivong A, Butsarakamruha T, Langlais RP. Antifungal activity of *Morinda citrifolia* fruit extract against *Candida albicans*. *Oral*

- Surg Oral Med Oral Pathol Oral Radiol Endod.* 2009;**108**(3):394–8. doi: [10.1016/j.tripleo.2009.05.044](https://doi.org/10.1016/j.tripleo.2009.05.044). [PubMed: [19716507](https://pubmed.ncbi.nlm.nih.gov/19716507/)].
15. Vermani K, Garg S. Herbal medicines for sexually transmitted diseases and AIDS. *J Ethnopharmacol.* 2002;**80**(1):49–66. doi: [10.1016/S0378-8741\(02\)00009-0](https://doi.org/10.1016/S0378-8741(02)00009-0).
 16. Grossman JH, Galask RP. Persistent vaginitis caused by metronidazole-resistant trichomonas. *Obstet Gynecol.* 1990;**76**(3-2):521–2.
 17. Burt S. Essential oils: Their antibacterial properties and potential applications in foods - A review. *Int J Food Microbiol.* 2004;**94**(3):223–53. doi: [10.1016/j.ijfoodmicro.2004.03.022](https://doi.org/10.1016/j.ijfoodmicro.2004.03.022). [PubMed: [15246235](https://pubmed.ncbi.nlm.nih.gov/15246235/)].
 18. Azadbakht M, Ziai H, Abdollahi F, Shabankhani B. [Effect of essential oils of Artemisia. Zataria and Myrtus on Trichomonas vaginalis]. *J Med Plant.* 2003;**4**(8):35–40. Persian.
 19. Ziaei H, Azadbakht M, Abdollahi F, Sha-bankhani B. [Effect of methanolic extracts of Artemisia aucheri Boiss, Zataria multiflora Boiss and Myrtus communis L. on Trichomonas vaginalis in vitro]. *J Gorgan Univ Med Sci.* 2006;**8**(1):34–8. Persian.
 20. Brandelli CL, Vieira Pde B, Macedo AJ, Tasca T. Remarkable anti-trichomonas vaginalis activity of plants traditionally used by the Mbya-Guarani indigenous group in Brazil. *Biomed Res Int.* 2013;**2013**:826370. doi: [10.1155/2013/826370](https://doi.org/10.1155/2013/826370). [PubMed: [23865068](https://pubmed.ncbi.nlm.nih.gov/23865068/)]. [PubMed Central: [PMC3705823](https://pubmed.ncbi.nlm.nih.gov/PMC3705823/)].
 21. Calzada F, Yopez-Mulia L, Tapia-Contreras A. Effect of Mexican medicinal plant used to treat trichomoniasis on Trichomonas vaginalis trophozoites. *J Ethnopharmacol.* 2007;**113**(2):248–51. doi: [10.1016/j.jep.2007.06.001](https://doi.org/10.1016/j.jep.2007.06.001). [PubMed: [17628366](https://pubmed.ncbi.nlm.nih.gov/17628366/)].
 22. Desrivot J, Waikedre J, Cabalion P, Herrenknecht C, Bories C, Hocquemiller R, et al. Antiparasitic activity of some New Caledonian medicinal plants. *J Ethnopharmacol.* 2007;**112**(1):7–12. doi: [10.1016/j.jep.2007.01.026](https://doi.org/10.1016/j.jep.2007.01.026). [PubMed: [17329051](https://pubmed.ncbi.nlm.nih.gov/17329051/)].
 23. Hassani S, Asghari G, Yousefi H, Kazemian A, Raffieian M, Darani HY. Effects of different extracts of Eucalyptus camaldulensis on Trichomonas vaginalis parasite in culture medium. *Adv Biomed Res.* 2013;**2**:47. doi: [10.4103/2277-9175.114187](https://doi.org/10.4103/2277-9175.114187). [PubMed: [24516847](https://pubmed.ncbi.nlm.nih.gov/24516847/)]. [PubMed Central: [PMC3905333](https://pubmed.ncbi.nlm.nih.gov/PMC3905333/)].
 24. Giordani RB, Vieira Pde B, Weizenmann M, Rosemberg DB, Souza AP, Bonorino C, et al. Candimine-induced cell death of the amitochondriate parasite Trichomonas vaginalis. *J Nat Prod.* 2010;**73**(12):2019–23. doi: [10.1021/np100449g](https://doi.org/10.1021/np100449g). [PubMed: [21105684](https://pubmed.ncbi.nlm.nih.gov/21105684/)].
 25. Vieira Pde B, Giordani RB, De Carli GA, Zuanazzi JA, Tasca T. Screening and bioguided fractionation of Amaryllidaceae species with anti-Trichomonas vaginalis activity. *Planta Med.* 2011;**77**(10):1054–9. doi: [10.1055/s-0030-1270740](https://doi.org/10.1055/s-0030-1270740). [PubMed: [21308614](https://pubmed.ncbi.nlm.nih.gov/21308614/)].
 26. Wang W, Wu N, Zu YG, Fu YJ. Antioxidative activity of Rosmarinus officinalis L. essential oil compared to its main components. *Food Chem.* 2008;**108**(3):1019–22. doi: [10.1016/j.foodchem.2007.11.046](https://doi.org/10.1016/j.foodchem.2007.11.046). [PubMed: [26065766](https://pubmed.ncbi.nlm.nih.gov/26065766/)].
 27. Nejati H, Farahpour MR, Neiriz Noghadeh M. [Experimental study of local antifungal activity of rosemary essential oil on Candida albicans infected cutaneous ulcer in rat: A study of cell pathology]. *Comp Pathobiol.* 2014;**11**(3):1397–405. Persian.
 28. Natanzian Ghahfarkhi M, Sattari M, Yadegari MH, Goudarzi GR, Sahrkhiz MJ. [Antifungal activity of essential oil and alcoholic extract of Carum copticum against fluconazole-resistant and susceptible Candida albicans isolated]. *Pathobiol Res.* 2008;**11**. Persian.
 29. Jafari AA, Jafari H, Deghanbanadkoki A, Baghbanian M. [Antifungal activity of aqueous extracts from Ferula assa foetida aerial parts on Candida albicans and its comparison with fluconazole in vitro]. *Toloo-e Behdasht.* 2014;**13**(3):171–81. Persian.
 30. Doddanna SJ, Patel S, Sundarrao MA, Veerabhadrapa RS. Antimicrobial activity of plant extracts on Candida albicans: An in vitro study. *Indian J Dent Res.* 2013;**24**(4):401–5. doi: [10.4103/0970-9290.118358](https://doi.org/10.4103/0970-9290.118358). [PubMed: [24047829](https://pubmed.ncbi.nlm.nih.gov/24047829/)].
 31. Prabhakar K, Kumar LS, Rajendran S, Chandrasekaran M, Bhaskar K, Sajit Khan AK. Antifungal activity of plant extracts against Candida species from oral lesions. *Indian J Pharm Sci.* 2008;**70**(6):801–3. doi: [10.4103/0250-474X.49128](https://doi.org/10.4103/0250-474X.49128). [PubMed: [21369447](https://pubmed.ncbi.nlm.nih.gov/21369447/)]. [PubMed Central: [PMC3040880](https://pubmed.ncbi.nlm.nih.gov/PMC3040880/)].
 32. Masomi F, Hassanshahian M. Antimicrobial activity of five medicinal plants on Candida albicans. *Iran J Toxicol.* 2016;**10**(6):39–43. doi: [10.29252/arakmu.10.6.39](https://doi.org/10.29252/arakmu.10.6.39).
 33. Bozin B, Mimica-Dukic N, Samojlik I, Jovin E. Antimicrobial and antioxidant properties of rosemary and sage (Rosmarinus officinalis L. and Salvia officinalis L., Lamiaceae) essential oils. *J Agric Food Chem.* 2007;**55**(19):7879–85. doi: [10.1021/jf0715323](https://doi.org/10.1021/jf0715323). [PubMed: [17708648](https://pubmed.ncbi.nlm.nih.gov/17708648/)].